

Perforation

Group by IBD type	Study name	Statistics for each study			Event rate and 95% CI					
		Event rate	Lower limit	Upper limit						Re
UC	lacopini, 2015	0.050	0.003	0.475	⊢-	-	— Ⅰ			
JC	Kasuga, 2021	0.045	0.003	0.448		_	_			
JC	Kinoshita, 2019	0.040	0.006	0.235						
IC	Matsumoto, 2019	0.042	0.003	0.425	├-	_	– I			
JC	Manta, 2021	0.057	0.018	0.161	-■-	-				
JC	Matsui, 2021	0.029	0.002	0.336	├-					
JC	Nishio, 2020	0.103	0.039	0.243	-■	$\boldsymbol{\vdash}$				
IC	Suzuki, 2017	0.016	0.001	0.206		— I				
IC	Yang DH, 2019	0.033	0.002	0.366	├-	-				
IC	Pooled	0.059	0.033	0.103						
IC	Prediction Interval				l i					
C+CD	Kochhar, 2018	0.071	0.004	0.577	┝╼	_				
IC+CD	Lightner, 2021	0.040	0.006	0.235	⊢					
JC+CD	Ngamruengphong, 2022	0.022	0.003	0.142		-				
JC+CD	Pooled	0.035	0.010	0.115		1				
JC+CD	Prediction Interval				T.	1				
Overall	Pooled	0.053	0.031	0.089	•	1				
Overall	Prediction Interval				l i					
					0.00	0.25	0.50	0.75	1.00	

overall I2 = 0% UC = 0% UC+CD = 0%

Supplementary Fig. 4. Forest plot for perforation. CI, confidence interval; UC, ulcerative colitis; CD, Crohn's disease.