

Supplementary Table 1. Type of Fecal Occult Blood Test in Each Institution

Institution	Number of patients	Method	Cut-off
1	8	FIT	Quantitative (50 ng/mL)
2	37	FIT	Qualitative
3	23	FIT	Quantitative (100 ng/mL)
4	6	FIT	Qualitative
5	14	FIT	Quantitative (99 ng/mL)
6	12	Chemical FOBT (guaiac)	Qualitative
7	4	FIT	Quantitative (100 ng/mL)
8	13	FIT	Quantitative (30 ng/mL)
9	24	FIT	Qualitative
10	3	FIT	Quantitative (100 ng/mL)

FIT, fecal immunochemical test; FOBT, fecal occult blood test.

Supplementary Table 2. Location of Bleeding Lesions in Patients with Positive Fecal Occult Blood Test

Finding of CE (<i>n</i>)	Duodenum	Jejunum	Ileum	Jejunum-ileum
Negative (11)	-	-	-	-
Angiodysplasia (10)	0	5	3	2
Bleeding of unidentified origin (2)	-	-	-	-
Non-specific erosion (1)	0	0	1	0
Non-specific ulcer (2)	0	1	0	1
Drug induced enteritis (1)	0	1	0	0
Crohn's disease (2)	0	0	2	0
Non-neoplastic polyp (3)	1	1	1	0
Submucosal tumor (2)	0	0	0	2
Neoplastic tumor (1)	0	1	0	0

CE, capsule endoscopy.